

**SUPPLEMENTARY MATERIALS TO:
COMPUTATIONAL INVESTIGATIONS, HIRSHFELD SURFACE ANALYSIS,
INTERACTION ENERGY CALCULATIONS, AND ENERGY FRAMEWORK
CRYSTAL STRUCTURE OF METHYL 2-AMINO-5-HYDROXYBENZOATE**

A. Saeed¹, G. Shabir¹, P.A. Channar¹, U. Flörke², T. Hökelek³, M.F. Erben⁴

¹Department of Chemistry, Quaid-I-Azam University, Islamabad, Islamic Republic of Pakistan

E-mail: shabirg@yahoo.com

²Hacettepe University, Department of Physics, Beytepe-Ankara, Republic of Turkey

³Department Chemie, Fakultät für Naturwissenschaften, Universität Paderborn, Paderborn, Federal Republic of Germany

⁴CEQUINOR (UNLP, CONICET-CCT La Plata), Departamento de Química, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Buenos Aires, República Argentina




Fig. S1. Calculated [B3LYP/6-31G(d,p)] potential function for internal rotation around the C1-C7 bond in the ester group of the title compound (for atom numbering see Fig. 1).




Fig. S2. Calculated [B3LYP/6-31G(d,p)] potential function for internal rotation around the C3-O3 bond in the hydroxyl group of title compound (for atom numbering see Fig. 1).