ЖУРНАЛ СТРУКТУРНОЙ ХИМИИ

2021. Том 62, № 12 *Декабрь* *DOI: 10.26902/JSC_id84207*

SUPPLEMENTARY MATERIALS TO: A FLUORESCENT PROBE OF THE Zn(II) COMPLEX CONSTRUCTED BY TERPHENYL-3,2″,3″,5,5″,5‴-HEXACARBOXYLIC ACID AND 3,5-BIS(1-IMIDAZOLE)PYRIDINE

X. Tian[1], Y. Li[1], Y. Zhang[1], E. Gao[1,2]

[1]*Key Laboratory of Liaoning Inorganic Molecule-Based Chemical and Department of Coordination Chemistry, Shenyang University of Chemical Technology, Shenyang, People's Republic of China*
E-mail: zhangy621@syuct.edu.cn
[2]*School of Chemical Engineering, University of Science and Technology Liaoning, Anshan, People's Republic of China*
E-mail: enjungao@ustl.edu.cn




Fig.S1 FT-IR spectrum of Zn complex.

© Tian X., Li Y., Zhang Y., Gao E., 2021




Fig.S2 PXRD patterns of Zn complex and simulared Zn complex.




Fig.S3 PXRD patterns of Zn complex after treatment for 24h in water at room temperature.




Fig.S4 The TGA curve of Zn complex